

TECHNIQUES IN MOLECULAR BIOLOGY

Course: Molecular Biology (02022312)

Instructor: Dr. M A Srouf

Textbook:

Watson J, Baker TA, Bell SP, Gann A, Levine M, Losick R (2008).
Molecular Biology of the Gene, 6th ed. [Chap 21/ pp739-82](#)

Lec # 3-4

Wed 01.02.2012

Nucleic acid Probes

- Previously cloned genes
- Synthetic oligonucleotides or DNA/RNA fragment
(complementary to part or all of target sequence)
- Radioactive labeled (^{32}P -dNTP)
- Nonradioactive labeled
 - ▣ DiG-dUTP
 - ▣ Biotin-dUTP



Nucleic acid Probes

- **The two basic methods for labeling probes are:**
- End labeling using T4 polynucleotide kinase:
 - ▣ adds the γ -phosphate from ^{32}P -ATP to the 5'-OH group
- Incorporation of labeled nucleotides into probe via PCR:
 - ▣ labeled nucleotides are usually nucleotides modified with either a radioactive atoms or fluorescent moiety.
 - ▣ Radioactive nucleotides typically have ^{32}P incorporated in the α -phosphate of one of the 4 dNTPs

Labeled DNA Probes



➤ Enzyme-Linked Probes

- Biotin /streptavidin
- Digoxigenin-UTP/antibody
- Radioactive ^{32}P



Factors Affecting Hybridization

- ❖ Temperature
- ❖ Ionic strength
- ❖ Probe length
- ❖ Probe mismatch
- ❖ % GC



Other forms of hybridization

□ Dot/Slot Blot

- ▣ DNA samples are spotted directly on the membrane and No electrophoresis is needed


□ Fluorescent in situ hybridization (FISH)

- combine with microscopy to localize cells expressing gene

DNA Cloning

- DNA cloning: the ability to construct recombinant DNA molecules and maintain them in cells
- A variety of techniques, often referred to as *recombinant DNA technology* are used in DNA cloning
- Isolation of a large amount of a single pure DNA molecule facilitates analysis of that DNA molecule.

DNA Cloning



Definition of terms used in cloning

- **Recombinant DNA:** any DNA molecule composed of sequences derived from different sources
- **Vector:** autonomously replicating genetic element used to carry an insert DNA (or cDNA fragment) into host cell for the purpose of gene cloning
- The most common host used is *E. coli* (genetically modified)
- **cDNA:** DNA molecule copied from an mRNA by reverse transcription

Recombinant vectors:

- Recombinant vectors: autonomously-replicating DNA used to 'carry' and amplify foreign DNA within host cells
- ***E. coli* plasmids**
- Phage lambda
- Cosmids (phage lambda + plasmid)
- BAC- bacterial artificial chromosomes
- YAC- yeast artificial chromosomes

Bacterial plasmids

- Occur naturally in bacteria (plasmids also present in single-cell eukaryotes, e.g, yeast)
- Circular dsDNA
- Autonomous replication
- Extra-chromosomal elements
- 1-200 kb size range
- Present in multiple copies per cell
- Transmitted during conjugation
- Exist in parasitic or symbiotic relationship with host cell (e.g, Antibiotic resistance)

Class activity?

- Why bacterial genomes and plasmids exist in a circular configuration??

Engineered / Recombinant plasmids

- **Basic elements of recombinant plasmids:**
 - Small in size >1.2-3 kb
 - Origin of replication, specific DNA sequences of 50-100bp> (Ori)
 - Selection marker (antibiotic resistance gene)
 - Polylinker/ multiple cloning site
- ❖ Most vectors are derived from pUC or pBR322
- ❖ Cloning capacity ~ up to 10kb (also up to 20kb or larger is possible, but is difficult)

Basic components of an *E. coli* “cloning plasmid” (Vector DNA)

- Basic elements of a cloning vector:
 1. Origin of replication
 2. Selectable marker, usually an antibiotic resistance gene
 3. Polylinker, with one or more unique sites for REs




Figure 5-12
Molecular Cell Biology, Sixth Edition
© 2008 W. H. Freeman and Company

Cleavage of DNA by EcoRI




Figure 5-11
Molecular Cell Biology, Sixth Edition
© 2008 W.H. Freeman and Company

Ligation of restriction fragments with complementary sticky ends.

Ligation of sticky ends is more efficient than blunt ends.






Figure 5-12
Molecular Cell Biology, Sixth Edition
© 2008 W.H. Freeman and Company



DNA cloning in a plasmid vector permits amplification of a DNA fragment




Figure 5-14
Molecular Cell Biology, Sixth Edition
© 2008 W. H. Freeman and Company






Figure 5-14 part 1
Molecular Cell Biology, Sixth Edition
© 2008 W. H. Freeman and Company



Genetic Transformation of Prokaryotes

- Transformation: the process by which a host organism can take up DNA from its environment
- **Two methods are commonly used:**
 1. Chemical transformation
 - Usually involves CaCl_2 and heat shock
 2. Electroporation
 - Electric field mediated membrane permeabilization
 - 10-100 times more efficient than chemical approach

Electroporation



Recombinant vectors:


- Recombinant vectors: autonomously-replicating DNA used to 'carry' and amplify foreign DNA within host cells
- *E. coli* plasmids
- **Phage lambda**
- Cosmids (phage lambda + plasmid)
- BAC- bacterial artificial chromosomes
- YAC- yeast artificial chromosomes

Phage Lambda (λ phage)

- Infection of *E. coli* by λ phage is 1000-fold more efficient than plasmid transformation
- Many more clones of phage can be grown on a single plate
- Infection and lysis of cells > ~100 phages / cell > plaques
- Can be assembled *in vitro*
- Cloning capacity ~ 25 kb


Bacteriophage Lambda genome and packaging of phage DNA

(a) λ Phage genome



Some regions of the genome can be replaced by exogenous DNA (diagonal lines) or deleted (dotted) without affecting the ability of phage to infect host cells and assemble new virions.

Bacteriophage Lambda: *in vitro* assembly




Recombinant vectors:

- ❑ Recombinant vectors: autonomously-replicating DNA used to 'carry' and amplify foreign DNA within host cells
- ❑ *E. coli* plasmids
- ❑ Phage lambda
- ❑ **Cosmids (phage lambda + plasmid)**
- ❑ BAC- bacterial artificial chromosomes
- ❑ YAC- yeast artificial chromosomes

Cosmids

- Combine the properties of plasmids and λ phage
- Cloning capacity 35-45 kb
- Common cosmids: pLFR-5 (6kb): a plasmid with 2 cos sites
- Cosmids based on P1 phage (115 kb) can carry up to 85kb insert

Cosmid Cloning System



λ -cos sites inserted into a small plasmid

Target DNA ligated between two cosmid DNA molecules

Recombinant DNA packaged and *E. coli*
Infected as before

Can clone DNAs up to 45 kb